



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

8. If the equation is of the first degree with respect to p , and algebraic with respect to x and y , p will be possible for every point of the plane and there will be no singular solution. If the equation is algebraic and of a degree higher than the second with respect to p we may apply the usual condition for equal roots that is to say, the equations $\varphi(x, y, p) = 0$ and $\varphi'(x, y, p) = 0$, where $\varphi' = \frac{d\varphi}{dp}$ must be satisfied by a common value of p : hence eliminating p between these equations we have the condition expressed as a relation between x and y . For example, given

$$\varphi = p^3 - 4xyp + 8y^2 = 0,$$

then

$$\varphi' = 3p^2 - 4xy = 0;$$

eliminating p we find $y = 0$, and $27y = 4x^3$.

Each of these is a branch of the envelop of the complete primitive, and is a singular solution. The complete primitive in fact is $y = c(x - c)^2$, representing a series of parabolas which touch the axis of x and the cubical parabola $27y = 4x^3$.

SOME TRIGONOMETRIC SERIES.

BY PROF. D. TROWBRIDGE, WATERBURGH, N. Y.

$$1. \text{ TAKE the equations } \log(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4, \dots (1)$$

$$\log(1-x) = -x - \frac{1}{2}x^2 - \frac{1}{3}x^3 - \frac{1}{4}x^4, \dots (2)$$

$$\theta \sqrt{-1} = \varphi, \quad 2 \cos \theta = e^{\varphi} + e^{-\varphi}, \quad 2 \sqrt{-1} \sin \theta = e^{\varphi} - e^{-\varphi}, \dots (3)$$

$$\log 2 \cos \theta = \varphi + \log(1 + e^{-2\varphi}) = -\varphi + \log(1 + e^{2\varphi}), \dots (4)$$

$$\log 2 \sin \theta = -\log \sqrt{-1} + \varphi + \log(1 - e^{-2\varphi}) = \log \sqrt{-1} - \varphi + \log(1 - e^{2\varphi}), (5)$$

$$2 \log 2 \cos \theta = \log(1 + e^{2\varphi}) + \log(1 + e^{-2\varphi}), \dots (6)$$

$$2 \log 2 \sin \theta = \log(1 - e^{2\varphi}) + \log(1 - e^{-2\varphi}). \dots (7)$$

Developing (6) and (7) by (1) and (2), we have

$$2 \log 2 \cos \theta = e^{2\varphi} + e^{-2\varphi} - \frac{1}{2}(e^{4\varphi} + e^{-4\varphi}) + \frac{1}{3}(e^{6\varphi} + e^{-6\varphi}), \dots (8)$$

$$2 \log 2 \sin \theta = -(e^{2\varphi} + e^{-2\varphi}) - \frac{1}{2}(e^{4\varphi} + e^{-4\varphi}) - \frac{1}{3}(e^{6\varphi} + e^{-6\varphi}) \dots (9)$$

$$\text{Whence we have } \log 2 \cos \theta = \cos 2\theta - \frac{1}{2} \cos 4\theta + \frac{1}{3} \cos 6\theta - \dots (10)$$

$$\log(2 \sin \theta)^{-1} = \cos 2\theta + \frac{1}{2} \cos 4\theta + \frac{1}{3} \cos 6\theta + \dots (11)$$

By differentiating (10) we have

$$\tan \theta = 2[\sin 2\theta - \sin 4\theta + \sin 6\theta - \dots] \dots (12)$$

$$2. \text{ From (4) we have } 2\varphi = 2\theta\sqrt{-1} = \log(1 + e^{2\phi}) - \log(1 + e^{-2\phi}) \\ = (e^{2\phi} - e^{-2\phi}) - \frac{1}{2}(e^{4\phi} - e^{-4\phi}) + \dots (13)$$

$$\text{Whence we have } \theta = \sin 2\theta - \frac{1}{2}\sin 4\theta + \frac{1}{3}\sin 6\theta - \dots (14)$$

Multiply this by $d\theta$ and integrate, multiply again and integrate, and continue this process. Denote the arbitrary constants by C_2, C_4, C_6 , &c., noticing that when the sine occurs in the series, this constant is zero. We shall have, if 1.2.3. . . . $n = [n]$,

$$\frac{\theta^2}{[2]} + C_2 = -\frac{1}{2} \left[\cos 2\theta - \frac{\cos 4\theta}{2^2} + \frac{\cos 6\theta}{3^2} \dots \right], \\ \frac{\theta^4}{[4]} + \frac{C_2\theta^2}{[2]} + C_4 = \frac{1}{2^3} \left[\cos 2\theta - \frac{\cos 4\theta}{2^4} + \frac{\cos 6\theta}{3^4} \dots \right], \\ \frac{\theta^n}{[n]} + \frac{C_2\theta^{n-2}}{[n-2]} + \frac{C_4\theta^{n-4}}{[n-4]} + \dots + C_n = \frac{(-1)^{\frac{n}{2}}}{2^{n-1}} \left[\cos 2\theta - \frac{\cos 4\theta}{2^n} + \frac{\cos 6\theta}{3^n} \dots \right] (15)$$

$$\text{Now make } S_n = 1 - \frac{1}{2^n} + \frac{1}{3^n} - \frac{1}{4^n} + \dots (16), S'_n = 1 + \frac{1}{2^n} + \frac{1}{3^n} + \frac{1}{4^n} \dots (17)$$

$$S''_n = 1 + \frac{1}{3^n} + \frac{1}{5^n} + \frac{1}{7^n} \dots (18), S'_n - S_n = \frac{1}{2^{n-1}} \left[1 + \frac{1}{2^n} + \frac{1}{3^n} \dots \right] = \frac{1}{2^{n-1}} S'_n;$$

$$S'_n = S_n \cdot \frac{2^{n-1}}{2^{n-1} - 1} \dots (19), S''_n = \frac{1}{2} (S'_n + S_n) = S_n \cdot \frac{2^n - 1}{2^{n-1} - 2} \dots (20)$$

Now make $\theta = \frac{1}{4}\pi = p$ in (15), and also make $\theta = 0$, and we find

$$\frac{p^n}{[n]} + \frac{C_2 p^{n-2}}{[n-2]} + \frac{C_4 p^{n-4}}{[n-4]} + \dots + C_n = \frac{(-1)^{\frac{n}{2}}}{2^{2^{n-1}}} S_n \dots (21); C_n = \frac{(-1)^{\frac{n}{2}}}{2^{n-1}} S_n \dots (22)$$

In these equations n is even. In (22) make $n = 2, 4, 6$, &c., and substitute in (21) and we have

$$\frac{p^n}{[n]} - \frac{S_2 p^{n-2}}{2.[n-2]} + \frac{S_4 p^{n-4}}{2^3[n-4]} - \frac{S_6 p^{n-6}}{2^5[n-6]} + \dots + \frac{(-1)^{\frac{n}{2}}}{2^{n-1}} S_n = \frac{(-1)^{\frac{n}{2}}}{2^{2^{n-1}}} S_n \dots (23)$$

From the form of (23) we easily see that p^n is a factor of all its terms; let us therefore make $\frac{1}{2^{n-1}} S_n = p^n B_n$, and we shall have

$$1 - n(n-1)B_2 + n(n-1)(n-2)(n-3)B_4 \dots + (-1)^{\frac{n}{2}} [n] B_n \cdot \frac{2^{n-1}}{2^n} = 0 \dots (24)$$

Now let $n = 2, 4, 6$, &c., in succession, and we shall have

$$B_2 = \frac{2}{3}, B_4 = \frac{1}{4 \cdot 5}, B_6 = \frac{1}{5 \cdot 3 \cdot 5}, \text{ \&c. } \dots \dots \dots (25)$$

$$\text{We also have } S_n = 2^{n-1} p^n B_n = \frac{1}{2^{n+1}} \pi^n B_n, S'_n = \frac{\pi^n B_n}{2^{n+1} - 4} \dots (26)$$

$$\text{Whence we find } S_2 = \frac{1}{12} \pi^2, S'_2 = \frac{1}{6} \pi^2, S''_2 = \frac{1}{8} \pi^2; \\ S_4 = \frac{7}{720} \pi^4, S'_4 = \frac{1}{90} \pi^4, S''_4 = \frac{1}{96} \pi^4$$

We have the following well-known relation :

$$\log \cos \theta = \log \left(1 - \frac{2^2 \theta^2}{\pi^2}\right) + \log \left(1 - \frac{2^2 \theta^2}{3^2 \pi^2}\right) + \log \left(1 - \frac{2^2 \theta^2}{5^2 \pi^2}\right) \dots \quad (27)$$

Developing by (2) we have

$$\begin{aligned} \log \cos \theta = & -\frac{2^2 \theta^2}{\pi^2} \left(1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots\right) - \frac{2^4 \theta^4}{2\pi^2} \left(1 + \frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + \dots\right) \\ & - \frac{2^6 \theta^6}{3\pi^6} \left(1 + \frac{1}{3^6} + \frac{1}{5^6} + \frac{1}{7^6} + \dots\right) \dots = -\frac{2^2 \theta^2}{\pi^2} S_2'' - \frac{2^4 \theta^4}{2\pi^4} S_4'' - \frac{2^6 \theta^6}{3\pi^6} S_6'' \dots \quad (28) \end{aligned}$$

If we differentiate (28) we shall have

$$\tan \theta = \frac{2^3 \theta}{\pi^2} S_2'' + \frac{2^5 \theta^3}{\pi^4} S_4'' + \frac{2^7 \theta^5}{\pi^6} S_6'' + \frac{2^9 \theta^7}{\pi^8} S_8'' + \dots; \quad \dots \quad (29)$$

$$\text{or } 2 \tan \theta = \frac{2^2 - 1}{2} B_2 \theta + \frac{2^4 - 1}{2^3 - 1} B_4 \theta^3 + \frac{2^6 - 1}{2^5 - 1} B_6 \theta^5 + \frac{2^8 - 1}{2^7 - 1} B_8 \theta^7 + \dots \quad (30)$$

3. Let us resume Eq. (12) and make

$$\tan \psi = 2[\sin 2\psi - \sin 4\psi + \sin 6\psi - \dots] = A_1 \psi + A_2 \psi^2 + A_3 \psi^3 + A_4 \psi^4 \dots \quad (31)$$

If we differentiate and make $\psi = 0$ after each differentiation, we shall find

$$\begin{aligned} A_1 &= 2^2[1 - 2 + 3 - 4 + \dots], \quad A_2 = 0, \quad A_4 = 0, \quad \&c. \\ [3] A_3 &= -2^4[1 - 2^3 + 3^3 - 4^3 + \dots], \\ [5] A_5 &= 2^6[1 - 2^5 + 3^5 - 4^5 + \dots], \quad \&c. \end{aligned}$$

Now make $1 - 2^{2n-1} + 3^{2n-1} - 4^{2n-1} + \dots = (-1)^{n+1} \Sigma(1)^{2n-1} \dots \dots \quad (32)$
then we shall have

$$\tan \psi = 2^2 \Sigma(1) \cdot \psi + \frac{2^4 \Sigma(1)^3}{[3]} \psi^3 + \frac{2^6 \Sigma(1)^5}{[5]} \psi^5 + \dots \dots \dots \quad (33)$$

We must now find the various sums represented by Σ . If we differentiate (14) we have $1 = 2[\cos 2\psi - \cos 4\psi + \cos 6\psi - \dots]$

$$0 = \sin 2\psi - 2 \sin 4\psi + 3 \sin 6\psi - \dots$$

$$0 = \cos 2\psi - 2^2 \cos 4\psi + 3^2 \cos 6\psi - \dots \quad \&c.$$

Make $\psi = 0$ and we see that $0 = 1 - 2^{2n} + 3^{2n} - 4^{2n} + \dots \dots \dots \quad (34)$

Now let $u = (1 + e^x)^{-1} = 1 - e^x + e^{2x} - e^{3x} + \dots, \dots \dots \quad (35)$
 e being the Napierian base of logarithms. If we differentiate (35) and make

$\frac{du}{dx} = u_1, \quad \frac{d^2 u}{dx^2} = u_2, \quad \frac{d^3 u}{dx^3} = u_3, \quad \&c.$ and retain the same notation when we make $x = 0$, we shall have $u = \frac{1}{2}, \quad u_1 = -\Sigma(1) = -\frac{1}{4}, \quad u_2 = -\Sigma(1)^2 = 0$, by (34), $u_3 = \Sigma(1)^3, \quad u_4 = 0, \quad u_5 = -\Sigma(1)^5, \quad \&c.$

From (35) we have $ue^x + u = 1. \quad \therefore e^x(u + u_1) + u_1 = 0, \quad e^x(u + 2u_1 + u_2) + u_2 = 0$, and generally

$$e^x \left[u + nu_1 + \frac{n(n-1)}{[2]} u_2 + \frac{n(n-1)(n-2)}{[3]} u_3 + \dots + nu_{n-1} + u_n \right] + u_n = 0$$

If we make $x = 0$, and remember that $u_2, u_4, \&c., = 0$, we shall have

$$u + nu_1 + \frac{n(n-1)(n-2)}{[3]}u_3 + \dots + 2u_n = 0, \dots \dots \dots (36)$$

in which n is an odd number. Or we may make

$$\frac{1}{2} = n\Sigma(1) - \frac{n(n-1)(n-3)}{[3]}\Sigma(1)^3 + \dots + (-1)^{\frac{n-1}{2}}2\Sigma(1)^n \dots (37)$$

If we now make $n = 1, 3, 5$, &c., in succession, we shall have

$$\Sigma(1) = \frac{1}{4}, \Sigma(1)^3 = \frac{1}{8}, \Sigma(1)^5 = \frac{1}{4}, \Sigma(1)^7 = \frac{1}{16}, \Sigma(1)^9 = \frac{3}{4}, \Sigma(1)^{11} = \frac{6}{8}, \dots \dots (38)$$

These values in (33) give

$$\tan \phi = \phi + \frac{2^4}{[3]}\frac{1}{8}\phi^3 + \frac{2^6}{[5]}\frac{1}{4}\phi^5 + \frac{2^8}{[7]}\frac{1}{16}\phi^7 + \frac{2^{10}}{[9]}\frac{3}{4}\phi^9 + \frac{2^{12}}{[11]}\frac{6}{8}\phi^{11} + \dots (39)$$

By comparing (30) and (33) we see that

$$\frac{2^2-1}{2-1}B_2 = 2, \frac{2^4-1}{2^3-1}B_4 = \frac{2^5}{[3]}\Sigma(1)^3, \dots \frac{2^{2n}-1}{2^{2n-1}-1}B_{2n} = \frac{2^{2n+1}}{[2n-1]}\Sigma(1)^{2n-1}. \dots (40)$$

We readily see that $u_1 = u^2 - u$, and hence

$$u_2 = (2u-1)u_1 = u(u-1)(2u-1) = 2u^3 - 3u^2 + u, \text{ and generally } u_n = A_{n+1}u^{n-1} - A_nu^n + A_{n-1}u^{n-1} - \dots + (-1)^nu, \dots \dots (41)$$

$$u_{n+1} = [(n+1)A_{n+1}u^n - nA_nu^{n-1} + (n-1)A_{n-1}u^{n-2} - \dots + (-1)^n](u^2 - u) = (n+1)A_{n+1}u^{n+2} - [nA_n + (n+1)A_{n+1}]u^{n+1} + [(n+1)A_{n-1} + nA_n]u^n - \dots + (-1)^{n+1}u. \dots (42)$$

By means of (41) and (42) we can easily compute the values of u_2, u_3 , &c., knowing the value of u_1 . We see that the sum of the coefficients of (42) is 0.

Since $2u = 1$, we may make $2u = z = 1$, and then

$$2^{n+1}u_n = A_{n+1}z^{n+1} - 2A_nz^n + 2^2A_{n-1}z^{n-1} - \dots + 2^n(-1)^nz, \dots \dots (43)$$

$$2^{n+2}u_{n+1} = (n+1)A_{n+1}z^{n+2} - 2[nA_n + (n+1)A_{n+1}]z^{n+1} + \dots + 2^{n+1}(-1)^{n+1}z \dots \dots (44)$$

Now let us make $2^{p+1}A_{n-p} = V_{n-p} \dots (45)$, then, if $2^{n+1}u_n = z_n \dots (46)$

$$\text{we shall have } z_n = V_{n+1}z^{n+1} - V_nz^n + V_{n-1}z^{n-1} - \dots + 2^n(-1)^nz, \dots (47)$$

$$z_{n+1} = (n+1)V_{n+1}z^{n+2} - [nV_n + 2(n+1)V_{n+1}]z^{n+1} + [(n-1)V_{n-1} + 2nV_n]z^n - \dots + 2^{n+1}(-1)^{n+1}z. \dots (48)$$

$$\text{Since } z = 1, z_n = V_{n+1} - V_n + V_{n-1} - \dots + 2^n(-1)^n, \dots \dots (49)$$

$$z_{n+1} = -(n+1)V_{n+1} + nV_n - (n-1)V_{n-1} + \dots + 2^n(-1)^{n+1} \dots \dots (50)$$

If we now make $n = 1$, we have $V_2 = 1, V_1 = 2$, and

$$\begin{aligned} z_1 &= z^2 - 2z &= 1 - 2 &= -1, \\ z_2 &= 2z^3 - 6z^2 + 4z &= -2 + 2 &= 0, \\ z_3 &= 6z^4 - 24z^3 + 28z^2 - 8z &= -6 + 12 - 4 &= 2, \\ z_4 &= 24z^5 - 120z^4 + 200z^3 - 120z^2 + 16z &= -24 + 72 - 56 + 8 &= 0. \end{aligned}$$

These equations are easily formed. We find $u_1 = \frac{1}{4}z_1 = -\frac{1}{2}$, $u_3 = \frac{1}{16}z_3 = \frac{1}{8}$, &c. It will be noticed that $\Sigma(1)^{2n+1}$ has some power of 2 for a denominator, and the relation of these numbers to the *Numbers of Bernoulli*, is easily ascertained.

If again we make $1.2.3 \dots n = [n]$, I have found

$$z_n = [n]z^{n+1} - [n+1]z^n + \frac{1}{6}[n+1](3n-2)z^{n-1} - \frac{1}{6}[n+1](n-1)(n-2)z^{n-2} \\ + \frac{1}{360}[n+1](15n^3-105n^2+230n-152)z^{n-3} - \dots \quad (51)$$

The remaining coefficients I have not yet obtained.

If we take

$$1 + x + x^2 + x^3 + \dots + x^n = \frac{x^{n+1} - 1}{x-1},$$

and make $x - 1 = y$, then $dx = dy$ and

$$1 + x + x^2 + \dots + x^n = \frac{(1+y)^{n+1} - 1}{y} = n + 1 + \frac{n(n+1)}{1.2}y \\ + \frac{n(n+1)(n-1)}{1.2.3}y^2 + \dots (1)$$

If we differentiate and make $\frac{n(n+1)}{1.2} = N_1$, $\frac{n(n+1)(n-1)}{1.2.3} = N_2$, &c.,

we shall have

$$1 + 2x + 3x^2 + \dots + nx^{n-1} = N_1 + 2N_2y + 3N_3y^2 + \dots \dots (2)$$

Multiply by $x = 1+y$, and make $A_1 = N_1 + 2N_2$, $A_2 = 2N_2 + 3N_3$, &c., and we shall have

$$x + 2x^2 + 3x^3 + \dots + nx^n = N_1 + A_1y + A_2y^2 + \dots \dots (3)$$

Differentiate again and we have

$$1 + 2^2x + 3^2x^2 + \dots + n^2x^{n-1} = A_1 + 2A_2y + 3A_3y^2 + \dots$$

Again multiply by $x = 1+y$, put $A_1 + 2A_2 = B_1$, $2A_2 + 3A_3 = B_2$, $3A_3 + 4A_4 = B_3$, &c., and continue the process, and then make $x = 1$, and $y = 0$ in these equations, and we shall have, if

$$S_n^{(p)} = 1 + 2^p + 3^p + \dots + n^p,$$

$$S_n^{(1)} = N_1,$$

$$S_n^{(2)} = A_1 = N_1 + 2N_2,$$

$$S_n^{(3)} = B_1 = A_1 + 2A_2,$$

$$S_n^{(4)} = C_1 = B_1 + 2B_2,$$

$$\&c. \quad \&c. \quad \&c.$$

$$A_2 = 2N_2 + 3N_3,$$

$$B_2 = 2A_2 + 3A_3,$$

$$\&c. \quad \&c.$$

$$A_3 = 3N_3 + 4N_4,$$

$$\&c. \quad \&c.$$